

# Existence and Attractors of Solutions for Nonlinear Parabolic Systems

By **A. El Hachimi and H. El Ouardi**

**Abstract:** We prove existence and asymptotic behaviour results for weak solutions of a mixed problem (S). We also obtain the existence of the global attractor and the regularity for this attractor in  $[H^2(\Omega)]^2$  and we derive estimates of its Hausdorff and fractal dimensions.

**Keywords:** Nonlinear parabolic systems; existence of solutions; global attractor; asymptotic behaviour; Hausdorff and fractal dimensions.

AMS subject classifications : 35K55, 35K57, 35K65, 35B40

## 0. Introduction

We consider the following nonlinear system

$$(S) \begin{cases} \frac{\partial b_1(u_1)}{\partial t} - \Delta u_1 + f_1(x, u_1, u_2) = 0 & \text{in } \Omega \times (0, T) \\ \frac{\partial b_2(u_2)}{\partial t} - \Delta u_2 + f_2(x, u_1, u_2) = 0 & \text{in } \Omega \times (0, T) \\ u_1 = u_2 = 0 & \text{in } \partial\Omega \times (0, T) \\ (b_1(u_1(x, 0)), b_2(u_2(x, 0))) = (b_1(\varphi_0(x)), b_2(\psi_0(x))) & \text{in } \Omega \end{cases}$$

where  $\Omega$  is a bounded open subset in  $\mathbb{R}^N$ ,  $N \geq 1$ , with a smooth boundary  $\partial\Omega$ . (S) is an example of nonlinear parabolic systems modelling a reaction diffusion process for which many results on existence, uniqueness and regularity have been obtained in the case where  $b_i(s) = s$  ( see, for instance [6, 7, 18]).

The case of a single equation of the type (S) is studied in [1, 2, 3, 4, 5, 8, 9, 19]. The purpose of this paper is the natural extension to system (S) of the results by [8], which concerns the single equation  $\frac{\partial \beta(u)}{\partial t} - \Delta u + f(x, t, u) = 0$ .

Actually, our work generalizes the question of existence and regularity of the global attractor obtained therein.

In the first section of this paper, we give some assumptions and preliminaries and in section 2, we prove the existence of absorbing sets and the existence of the global attractor; while in section 3, we present the regularity of the attractor and show stabilization property. Finally, section 4 is devoted to estimates of the Hausdorff and fractal dimensions.

## 1. Preliminaries, Existence and Uniqueness

### 1.1 Notations and Assumptions

Let  $b_i$ , ( $i = 1, 2$ ) be continuous functions with  $b_i(0) = 0$ . We define for  $t \in \mathbb{R}$

$\Psi_i(t) = \int_0^t b_i(\tau) d\tau$ . Then the Legendre transform  $\Psi_i^*$  of  $\Psi_i$  is defined by  $\Psi_i^*(\tau) = \sup_{s \in \mathbb{R}} \{\tau s - \Psi_i(s)\}$ .  $\Omega$  stands for a regular open bounded subset of

$R^N$  and for any  $T > 0$ , we set  $Q_T = \Omega \times (0, T)$  and  $S_T = \partial\Omega \times (0, T)$ , where  $\partial\Omega$  is the boundary of  $\Omega$ . The norm in a space  $X$  will be denoted by  $\|\cdot\|_r$  if  $X = L^r(\Omega)$  for all  $r : 1 \leq r \leq +\infty$ ,  $\|\cdot\|_X$  otherwise and  $\langle \cdot, \cdot \rangle_{X, X'}$  will denote the duality product between  $X$  and its dual  $X'$ .

We start by introducing our assumptions and making precise the meaning of a solution of (S). Consider the system (S) under the following assumptions:

(H1)  $(\varphi_0, \psi_0) \in L^2(\Omega) \times L^2(\Omega)$ .

(H2)  $b_i$  is an increasing continuous function from  $\mathbb{R}$  into  $\mathbb{R}$ ,  $b_i(0) = 0$ , and there exists  $c_{ij} > 0$  such that  $|b_i(s)| \leq c_{i1}|s| + c_{i2}$ , for all  $s \in \mathbb{R}$ ,  $i = 1, 2$ .

(H3)  $f_i \in C^1(\overline{\Omega} \times \mathbb{R} \times \mathbb{R})$ .

(H4)  $\forall x \in \Omega, \forall \xi \in \mathbb{R}, \exists c_1 > 0, c_2 > 0 :$

$$\begin{cases} \text{sign}(\xi)f_1(x, \xi, 0) \geq -c_1 \\ \text{sign}(\xi)f_2(x, 0, \xi) \geq -c_2. \end{cases}$$

(H5) For any  $N > 0, \exists c_3 > 0, c_4 > 0, c_5 > 0 :$

$$\begin{cases} \text{sign}(\xi)f_1(x, \xi, v) \geq c_3|\xi|^{p_1-1} - c_4 \\ |f_1(x, \xi, v)| \leq c_5(|\xi|^{p_1-1} + 1) & p_1 > 2 \\ |f(x, u, v)| \leq a_1(|u|), \text{ where } a : \mathbb{R}^+ \rightarrow \mathbb{R}^+ \text{ is increasing} \\ \text{for any } v : |v| \leq N. \end{cases}$$

(H6) For any  $M > 0, \exists c_6 > 0, c_7 > 0, c_8 > 0 :$

$$\begin{cases} \text{sign}(\xi)f_2(x, u, \xi) \geq c_6|\xi|^{p_2-1} - c_7 \\ |f_2(x, u, \xi)| \leq c_8(|\xi|^{p_2-1} + 1) & p_2 > 2 \\ |f_2(x, u, v)| \leq a_2(|v|), \text{ where } a_2 : \mathbb{R}^+ \rightarrow \mathbb{R}^+ \text{ is increasing} \\ \text{for any } u : |u| \leq M. \end{cases}$$

(H7)  $0 < \gamma_i \leq b'_i(s)$  for all  $s \in \mathbb{R}$ .

**Definition** By a weak solution of (S), we mean an element

$$u_i \in L^{p_i}(0, T; L^{p_i}(\Omega)) \cap L^2(0, T; H_0^1(\Omega)) \cap L^\infty(t_0, T; L^\infty(\Omega)), \text{ for all } t_0 > 0$$

such that

$$\frac{\partial b_i(u_i)}{\partial t} \in L^{p_i^*}(0, T; L^{p_i^*}(\Omega)) + L^2(0, T; H^{-1}(\Omega)) \text{ and } \forall \phi_i \in L^2(0, T; H^{-1}(\Omega)) :$$

$$\int_0^T \left\langle \frac{\partial b_i(u_i)}{\partial t}, \phi_i \right\rangle_{V_i^*, V_i} dt + \int_0^T \int_\Omega \nabla u_i \nabla \phi_i dx dt + \int_0^T \int_\Omega f_i(x, u_1, u_2) dx dt = 0,$$

and if  $(\phi_i)_t \in L^2(0, T; L^2(\Omega)), \phi_i(T) = 0$

$$\int_0^T \left\langle \frac{\partial b_i(u_i)}{\partial t}, \phi_i \right\rangle_{V_i', V_i} dt = - \int_0^T \int_\Omega (b_i(u_i(t)) - b_i(u_i(x, 0))) (\phi_i)_t dx dt,$$

where  $V_i = L^{p_i}(\Omega) \cap H_0^1(\Omega)$ ,  $V_i' = L^{p_i'}(\Omega) + H^{-1}(\Omega)$ ,  $\frac{1}{p_i} + \frac{1}{p_i'} = 1$ ,  $i = 1, 2$ .

## 1.2. Existence theorem.

**Theorem 1** Let (H1) to (H6) be satisfied. Then there exists a solution  $(u_1, u_2)$  of problem (S) such that for  $i = 1, 2$ , we have

$$u_i \in L^{p_i}(0, T; L^{p_i}(\Omega)) \cap L^2(0, T; H_0^1(\Omega)) \cap L^\infty(t_0, T; L^\infty(\Omega)), \forall t_0 > 0$$

Proof: By theorem 3.2 in [8], we can choose  $u_i^0 \in L^{p_i}(Q_T) \cap L^2(0, T; H_0^1(\Omega)) \cap L^\infty(\tau, T; L^\infty(\Omega))$ , for any  $\tau > 0$  such that :

$$\begin{cases} \frac{\partial b_1(u_1^0)}{\partial t} - \Delta u_1^0 + f_1(x, u_1^0, 0) = 0 & \text{in } Q_T \\ u_1^0 = 0 & \text{in } S_T \\ b_1(u_1^0)_{t=0} = b_1(\varphi_0) & \text{in } \Omega \end{cases}$$

$$\begin{cases} \frac{\partial b_2(u_2^0)}{\partial t} - \Delta u_2^0 + f_2(x, 0, u_2^0) = 0 & \text{in } Q_T \\ u_2^0 = 0 & \text{in } S_T \\ b_1(u_2^0)_{t=0} = b_1(\psi_0) & \text{in } \Omega \end{cases}$$

and we construct two sequences of functions  $(u_1^n)$  and  $(u_2^n)$ , such that :

$$\begin{cases} \frac{\partial b_1(u_1^n)}{\partial t} - \Delta u_1^n + f(x, u_1^n, u_2^{n-1}) = 0 & \text{in } Q_T \\ u_1^n = 0 & \text{in } S_T \end{cases} \quad (1.1)$$

$$\begin{cases} b_1(u_1^n)_{t=0} = b_1(\varphi_0) & \text{in } \Omega \end{cases} \quad (1.2)$$

$$\begin{cases} \frac{\partial b_2(u_2^n)}{\partial t} - \Delta u_2^n + f_2(x, u_1^{n-1}, u_2^n) = 0 & \text{in } Q_T \\ u_2^n = 0 & \text{in } S_T \end{cases} \quad (1.4)$$

$$\begin{cases} b_2(u_2^n)_{t=0} = b_2(\psi_0) & \text{in } \Omega \end{cases} \quad (1.5)$$

We need lemma 1 and lemma 2 below to complete the proof of theorem 1. From now on we denote by  $c_i$  various positive constants independent of  $n$ .

**Lemma 1**

$$\forall \tau > 0, \exists c_\tau > 0 \quad \text{such that} \quad \|u_i^n\|_{L^\infty(\tau, T; L^\infty(\Omega))} \leq c_\tau. \quad (1.7)$$

Proof : For  $n = 0$ , (1.7) is proved in [7]. So, suppose (1.7) for  $(n - 1)$ .

Multiplying (1.1) by  $|b_1(u_1^n)|^k b_1(u_1^n)$  and using (H2),(H5) , we obtain :

$$\frac{1}{k+2} \int_{\Omega} |b_1(u_1^n)|^{k+2} dx + c_9 \int_{\Omega} |b_1(u_1^n)|^{k+p_1} dx \leq c_{10} \int_{\Omega} |b_1(u_1^n)|^{k+1} dx.$$

Setting  $y_{k,n}(t) = \|b_1(u_1^n)\|_{L^{k+2}(\Omega)}$  and using Holder's inequality on both sides, we have the existence of two constants  $\lambda > 0$  and  $\delta > 0$  such that

$$\frac{dy_{k,n}(t)}{dt} + \lambda y_{k,n}^{p_1-1}(t) \leq \delta,$$

which implies from lemma 5.1([22]) that  $\forall t \geq \tau > 0$

$$y_{k,n}(t) \leq \left(\frac{\delta}{\lambda}\right)^{\frac{1}{p_1-1}} + \frac{1}{[\lambda(p_1 - 2)t]^{\frac{1}{p_1-2}}}.$$

As  $k \rightarrow \infty$ , we obtain

$$|u_1^n(t)| \leq c_\tau \quad \forall t \geq \tau > 0.$$

The same holds also for  $u_2^n$ . ■

**Lemma 2**  $\forall \tau > 0, \exists c_i = c_i(\tau, \varphi_0, \psi_0) > 0 :$

$$\|u_i^n\|_{L^2(0, T; H_0^1(\Omega))} \leq c_{11},$$

$$\|u_i^n\|_{L^\infty(\tau, T; H_0^1(\Omega) \cap L^\infty(\Omega))} \leq c_{12}$$

and

$$\sum_{i=1}^2 \left[ \int_0^T \int_{\Omega} |\nabla u_i^n|^2 dx + c_{13} \int_0^T \int_{\Omega} |u_i^n|^{p_i} dx \right] \leq c_{14}.$$

Proof of lemma 2 : Multiplying (1.1) by  $u_1^n$  and (1.4) by  $u_2^n$ , and adding , we get :

$$\frac{d}{dt} \sum_{i=1}^2 \left[ \int_{\Omega} \Psi_i^*(b_i(u_i^n)) dx \right] + \sum_{i=1}^2 \int_{\Omega} |\nabla u_i^n|^2 dx + c_{15} \sum_{i=1}^2 \int_{\Omega} |u_i^n|^{p_i} dx \leq c_{16}. \quad (1.8)$$

But

$$|\varphi_0|_{L^2(\Omega)} + |\psi_0|_{L^2(\Omega)} \leq c \Rightarrow \int_{\Omega} \Psi_1^*(b_1(\varphi_0)) dx + \int_{\Omega} \Psi_2^*(b_2(\psi_0)) dx \leq c,$$

$$\text{so we deduce that } \sum_{i=1}^2 \int_0^T \int_{\Omega} |\nabla u_i^n|^2 dx + c_{17} \sum_{i=1}^2 \int_0^T \int_{\Omega} |u_i^n|^{p_i} dx \leq c_{18}.$$

Whence lemma 2.

From lemma 2 and Lemma 1, there is a subsequence  $u_i^n$  ( $i = 1, 2$ ) with the following properties:

$$u_i^n \rightharpoonup u_i \text{ weakly in } L^2(0, T; H_0^1(\Omega)) \cap L^{p_i}(0, T; L^{p_i}(\Omega))$$

$$b_i(u_i^n) \rightharpoonup \chi_i \text{ weakly in } L^2(0, T; L^2(\Omega))$$

$b_i(u_i^n) \rightarrow \chi_i$  strongly in  $L^2(\tau, T; H^{-1}(\Omega))$  ( by the compactness result of Aubin ( see [22])). By lemma 7([9]), we have  $\chi_i = b_i(u_i)$ . Moreover,

$$f_1(., u_1^n, u_2^{n-1}) \text{ converges to } f_1(., u_1, u_2) \text{ in } L^r(\tau, T; L^r(\Omega)), \forall r \geq 1, \forall \tau \geq 1$$

$$\text{and } f_2(., u_1^{n-1}, u_2^n) \text{ converges to } f_2(., u_1, u_2) \text{ in } L^r(\tau, T; L^r(\Omega)), \forall r \geq 1;$$

taking the limit as  $n$  goes to  $\infty$  , we deduce that  $(u_1, u_2)$  is a weak solution of (S).

### 1.3. Uniqueness

#### Theorem 2.

Assume that  $f_1$  and  $f_2$  verify :

$$(H8) \begin{cases} \forall M > 0, \forall N > 0, \exists c_M > 0, c_N > 0 : \\ \forall u, \bar{u}, v, \bar{v} : |u| + |\bar{u}| \leq M \text{ and } |v| + |\bar{v}| \leq N, \text{ we have} \\ |f_1(x, u, v) - f_1(x, \bar{u}, \bar{v})|^2 + |f_2(x, u, v) - f_2(x, \bar{u}, \bar{v})|^2 \leq \\ c_M(b_1(u) - b_1(\bar{u}))(u - \bar{u}) + c_N(b_2(v) - b_2(\bar{v}))(v - \bar{v}). \end{cases}$$

Then (S) has a unique solutions  $(u, v)$  in  $Q_T$ .

Proof : Let  $(u, v)$  and  $(\bar{u}, \bar{v})$  be solutions of (S); then we have :

$$\frac{\partial(b_1(u) - b_1(\bar{u}))}{\partial t} - \Delta(u - \bar{u}) = f_1(x, \bar{u}, \bar{v}) - f_1(x, u, v) \quad (1.9)$$

and

$$\frac{\partial(b_2(v) - b_2(\bar{v}))}{\partial t} - \Delta(v - \bar{v}) = f_2(x, \bar{u}, \bar{v}) - f_2(x, u, v). \quad (1.10)$$

Multiplying (1.9) by  $w_1 = (-\Delta)^{-1}(b_1(u) - b_1(\bar{u}))$  and (1.10) by  $w_2 = (-\Delta)^{-1}(b_2(v) - b_2(\bar{v}))$  and adding, we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left[ \|b_1(u) - b_1(\bar{u})\|_{H^{-1}(\Omega)}^2 + \|b_2(v) - b_2(\bar{v})\|_{H^{-1}(\Omega)}^2 \right] + \\ & (b_1(u) - b_1(\bar{u}), u - \bar{u})_{L^2(\Omega)} + (b_2(v) - b_2(\bar{v}), v - \bar{v})_{L^2(\Omega)} \leq \\ & c \|f_1(x, u, v) - f_1(x, \bar{u}, \bar{v})\|_{H^{-1}(\Omega)} \|b_1(u) - b_1(\bar{u})\|_{H^{-1}(\Omega)} + \end{aligned}$$

$$c \|f_2(x, u, v) - f_2(x, \bar{u}, \bar{v})\|_{H^{-1}(\Omega)} \|b_2(v) - b_2(\bar{v})\|_{H^{-1}(\Omega)} . \quad (1.11)$$

From hypothesis (H8) we obtain

$$\begin{aligned} & \|f_1(x, u, v) - f_1(x, \bar{u}, \bar{v})\|_{H^{-1}(\Omega)}^2 + \|f_2(x, u, v) - f_2(x, \bar{u}, \bar{v})\|_{H^{-1}(\Omega)}^2 \\ & \leq c \left[ \|f_1(x, u, v) - f_1(x, \bar{u}, \bar{v})\|_{L^2(\Omega)}^2 + \|f_2(x, u, v) - f_2(x, \bar{u}, \bar{v})\|_{L^2(\Omega)}^2 \right] \\ & \leq c c_M (b_1(u) - b_1(\bar{u}), u - \bar{u})_{L^2(\Omega)} + c c_N (b_2(v) - b_2(\bar{v}), v - \bar{v})_{L^2(\Omega)} , \quad (1.12) \end{aligned}$$

where  $M = \|u\|_{L^\infty(0,T;L^\infty(\Omega))} + \|\bar{u}\|_{L^\infty(0,T;L^\infty(\Omega))}$  and  $N = \|v\|_{L^\infty(0,T;L^\infty(\Omega))} + \|\bar{v}\|_{L^\infty(0,T;L^\infty(\Omega))}$ .

Therefore, using Schwartz inequality in (1.11), the fact that  $(b_i, i = 1, 2)$  is increasing and (1.12), we deduce that

$$\begin{aligned} & \frac{d}{dt} \left[ \|b_1(u) - b_1(\bar{u})\|_{H^{-1}(\Omega)}^2 + \|b_2(v) - b_2(\bar{v})\|_{H^{-1}(\Omega)}^2 \right] \leq \\ & c \frac{d}{dt} \left[ \|b_1(u) - b_1(\bar{u})\|_{H^{-1}(\Omega)}^2 + \|b_2(v) - b_2(\bar{v})\|_{H^{-1}(\Omega)}^2 \right] . \end{aligned}$$

Thus, we deduce that  $b_1(u) = b_1(\bar{u})$  and  $b_2(v) = b_2(\bar{v})$ , hence  $u = \bar{u}$  and  $v = \bar{v}$ .

**Remark 1.** Theorem 1 establishes the existence of dynamical system  $\{S(t)\}_{t \geq 0}$  which maps  $L^2(\Omega) \times L^2(\Omega)$  into  $L^2(\Omega) \times L^2(\Omega)$  such that  $S(t)(\varphi_0, \psi_0) = (u_1(t), u_2(t))$ .

## 2. Global attractor

**Proposition 1** Assume that (H1)-(H8) hold; then the solution  $(u_1, u_2)$  of system (S) satisfies :

$$|u_1(t)|_{L^\infty(\Omega)} + |u_2(t)|_{L^\infty(\Omega)} \leq c(t_0) \quad \forall t \geq t_0 \quad (2.0)$$

and

$$\sum_{i=1}^2 \int_{\Omega} |\nabla u_i|^2 dx \leq c \quad \forall t \geq t_0 + r \quad (2.1)$$

Proof : Reasoning as the proof of lemma 1, we also have (2.0).

Multiplying the first equation of (S) by  $u_1$  and the second by  $u_2$ , by (H2) and (2.5), we get :

$$\frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \Psi_i^*(b_i(u_i)) dx + \sum_{i=1}^2 \frac{1}{2} \int_{\Omega} |\nabla u_i|^2 dx = - \sum_{i=1}^2 \int_{\Omega} f_i(x, u) u_i dx \leq c. \quad (2.2)$$

For fixed  $r > 0$  and  $\tau > 0$ , integrate (2.2) on  $]t, t + r[$

$$\forall t \geq \tau > 0 \quad \sum_{i=1}^2 \int_t^{t+r} \int_{\Omega} |\nabla u_i|^2 dx ds \leq c(\tau). \quad (2.3)$$

Multiplying the first equation of (S) by  $(u_1)_t$  and the second by  $(u_2)_t$ , we get

$$\frac{1}{2} \frac{d}{dt} \left( \sum_{i=1}^2 \int_{\Omega} |\nabla u_i|^2 dx \right) + \sum_{i=1}^2 \int_{\Omega} (b'_i(u_i)) \left( \frac{\partial u_i}{\partial t} \right)^2 dx = \quad (2.4)$$

$$\sum_{i=1}^2 \int_{\Omega} f_i(x, u) (u_i)_t \leq \frac{1}{2} \sum_{i=1}^2 \int_{\Omega} \frac{f_i^2(x, u)}{b'_i(u_i)} dx + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega} (b'_i(u_i)) \left( \frac{\partial u_i}{\partial t} \right)^2 dx.$$

By (H7) and the properties of functions  $f_i$ , we obtain:

$$\frac{d}{dt} \left[ \sum_{i=1}^2 \int_{\Omega} |\nabla u_i|^2 dx \right] \leq c(\tau). \quad (2.5)$$

From the uniform Gronwall's lemma see [22], we get (2.1). ■

**Remark 2.** By proposition 1 we deduce that there exist absorbing sets in  $L^{\sigma_1}(\Omega) \times L^{\sigma_1}(\Omega)$  for any  $\sigma_i : 1 \leq \sigma_i \leq +\infty$  and absorbing sets in  $H_0^1(\Omega) \times H_0^1(\Omega)$ ; then assumptions (1.1), (1.4) and (1.12) in theorem 1.1 [22, p.23] are satisfied with  $U = [L^2(\Omega)]^2$ , so we have the following :

**Theorem 2.** Assume that (H1) - (H7) are satisfied. Then the semi-group  $S(t)$  associated with the boundary value problem (S) possesses a maximal attractor  $A$ , which is bounded in  $[H_0^1(\Omega) \cap L^\infty(\Omega)] \times [H_0^1(\Omega) \cap L^\infty(\Omega)]$ , compact and connected in  $[L^2(\Omega)]^2$ . Its domain of attraction is the whole space  $[L^2(\Omega)]^2$ .

### 3. A regularity property of the attractor

In this section we shall show supplementary regularity estimates on the solution of problem (S) and by use of them, we shall obtain more regularity on the attractor obtained in section 3. We shall assume that

$$(H9) \{ N \leq 3 \text{ and } b_i \text{ is of class } \mathcal{C}^3.$$

Hereafter, we shall assume that there exist positive constants  $\delta_i > 0$  and a function  $\Phi$  from  $R^{N+2}$  to  $R$  such that :

$$(H10) \left\{ \begin{array}{l} f_i(x, u) = f_i(u) - h_i(x) = \delta_i \frac{\partial \Phi}{\partial u_i} \\ f_i \text{ satisfying (H3) to (H6) and } h_i \in L^\infty(\Omega). \end{array} \right.$$

We shall denote :  $r(t) = \sum_{i=1}^2 \int_{\Omega} b'_i(u_i) (u'_i)^2 dx$ .

**Theorem 3** Let  $f_i$  and  $b_i$  satisfies hypothesis (H1) to (H10). Then the solution  $(u_1, u_2)$  of problem (S) satisfies the following regularity estimates:

$$\frac{\partial b_i(u_i)}{\partial t} \in L^2(t_0, +\infty; L^2(\Omega)), \quad (3.0)$$

$$\frac{\partial \nabla u_i}{\partial t} \in L^2(t_0, +\infty; L^2(\Omega)), \quad (3.1)$$

and

$$u_i \in H^2(\Omega). \quad (3.2)$$

To prove this theorem, we need the following lemma:

**Lemma 3** Under the assumptions of theorem 3, there exist constants  $C = C(\varphi_0, \psi_0)$ , such that for any  $T > 0$  :

$$\|u_i\|_{L^\infty(0, T, H_0^1(\Omega))} \leq C < \infty \quad (3.3)$$

and

$$\left\| \frac{\partial u_i}{\partial t} \right\|_{L^2(Q_T)} \leq C < \infty. \quad (3.4)$$

**Proof of lemma 3 :** Multiplying the equation  $\frac{\partial b_i(u_i)}{\partial t} - \text{div} [\nabla u_i] + \delta_i \frac{\partial \Phi}{\partial u_i} = 0$  by  $\frac{1}{\delta_i} (u_i)_t$  and adding the two equations, we obtain :

$$\begin{aligned} & \sum_{i=1}^2 \frac{1}{\delta_i} \int_{Q_T} b'_i(u_i) \left( \frac{\partial u_i}{\partial t} \right)^2 dx dt + \sum_{i=1}^2 \frac{1}{2\delta_i} \int_{\Omega} |\nabla u_i(., T)|^2 dx = \\ & \int_{\Omega} [-\Phi(., u_1(T), u_2(T)) + \Phi(., \varphi_0, \psi_0)] dx = \frac{1}{2\delta_1} \int_{\Omega} |\nabla \varphi_0|^2 dx + \frac{1}{2\delta_2} \int_{\Omega} |\nabla \psi_0|^2 dx. \end{aligned} \quad (3.5)$$

$\Phi$  being continuous and  $(u_1, u_2)$  bounded, we then obtain:

$$\sum_{i=1}^2 \frac{\gamma_i}{\delta_i} \int_{Q_T} \left( \frac{\partial u_i}{\partial t} \right)^2 dx dt + \sum_{i=1}^2 \frac{1}{2\delta_i} \int_{\Omega} |\nabla u_i(., T)|^2 dx \leq C(\varphi_0, \psi_0), \quad (3.6)$$

whence (3.3) and (3.4). ■

**Proof of theorem 3 :** Differentiating equation  $\frac{\partial b_i(u_i)}{\partial t} - \operatorname{div} [\nabla u_i] + f_i(u_1, u_2) = h_i$  we obtain

$$b'_i(u_i)u''_i + b''_i(u_i)(u'_i)^2 - \operatorname{div} (\nabla u_i))' + \sum_{j=1}^2 \frac{\partial f_i(u)}{\partial u_j} u'_j = 0 . \quad (3.7)$$

Now multiplying (3.7) by  $u'_i$ , and integrating over  $\Omega$  gives

$$\frac{1}{2}r'(t) + \frac{1}{2} \sum_{i=1}^2 \int_{\Omega} b''_i(u_i)(u'_i)^3 dx + \sum_{i=1}^2 \|u'_i\|_{H_0^1(\Omega)} + \int_{\Omega} \left( \sum_{i=1}^2 \sum_{j=1}^2 \frac{\partial f_i(u)}{\partial u_j} u'_j \right) u'_i dx = 0 . \quad (3.8)$$

The  $L^\infty$  estimate and hypothesis ( H9) imply successively :

$$\int_{\Omega} \left( \sum_{i=1}^2 \sum_{j=1}^2 \frac{\partial f_i(u)}{\partial u_j} u'_j \right) u'_i dx \leq M \sum_{i=1}^2 \int_{\Omega} (u'_i)^2 dx, \quad (3.9)$$

$$\gamma \sum_{i=1}^2 \int_{\Omega} (u'_i)^2 dx \leq r(t) \leq M \sum_{i=1}^2 \|u'_i\|_{H_0^1(\Omega)}^2, \quad (3.10)$$

and

$$-\frac{1}{2} \sum_{i=1}^2 \int_{\Omega} b''(u_i)(u'_i)^3 dx \leq \sum_{i=1}^2 \frac{M_i}{2} |u'_i|_{L^3(\Omega)}^3 . \quad (3.11)$$

Since for  $N \leq 3$ ,  $H_0^1(\Omega)$  is continuously imbedded in  $L^6(\Omega)$ , we then obtain by Young's inequality that :

$$|u'_i|_{L^3(\Omega)}^3 \leq c |u'_i|_{L^2(\Omega)}^{\frac{9}{4}} |u'_i|_{L^3(\Omega)}^3 \leq c |u'_i|_{L^2(\Omega)}^{\frac{18}{5}} + \frac{1}{2} \|u'_i\|_{H_0^1(\Omega)}^2 . \quad (3.12)$$

By (3.9),(3.10),(3.11) and (3.12), (3.7) becomes :

$$r'(t) + \frac{1}{2} \sum_{i=1}^2 \|u'_i\|_{H_0^1(\Omega)}^2 + cr(t) \leq cr(t)^{\frac{9}{5}} + cr(t) \leq cr(t)^2 + c. \quad (3.13)$$

On the other hand, using (2.4) we obtain :

$$\sum_{i=1}^2 \int_{\tau}^{\tau+r} \int_{\Omega} b'_i(u_i) (u'_i)^2 dx dt \leq c_{\tau}, \text{ for any } \tau \geq t_0 . \quad (3.14)$$



Estimates (3.13) and (3.14) and the use of the uniform Gronwall' lemma thus gives

$$r(t) \leq c(t_0), \text{ for any } \forall t \geq t_0. \quad (3.15)$$

Now, by (3.15) and hypothesis (H1) , we get :

$$\sum_{i=1}^2 \int_{\Omega} \left( \frac{\partial b_i(u_i)}{\partial t} \right)^2 dx \leq Mr(t) \leq c(t_0) \text{ for any } t \geq t_0 .$$

Then (3.0) is satisfied. Now, as we have :

$$-\Delta u_i = -f_i(x, u) - b_i(u_i)_t \in L^\infty(t_0, +\infty; L^2(\Omega)),$$

then by (S)  $u_i(., t)$  is in bounded subset of  $H^2(\Omega)$ . Hence estimate (3.2) follows. ■

For a solution  $(u_1, u_2)$  of (S), we define the  $\omega$  - limit set by :

$$\omega(\varphi_0, \psi_0) = \left\{ \begin{array}{l} w = (w_1, w_2) \in (H_0^1(\Omega) \times L^\infty(\Omega)) \cap (H_0^1(\Omega) \times L^\infty(\Omega)) \\ \exists t_n \rightarrow +\infty \quad u_1(., t_n) \rightarrow w_1 \text{ in } L^2(\Omega), \quad u_2(., t_n) \rightarrow w_2 \text{ in } L^2(\Omega) \end{array} \right\}.$$

**Corollary 1.** Under the assumptions (H1) to (H10), we have  $\omega(\varphi_0, \psi_0) \neq \emptyset$  and any  $(w_1, w_2) \in \omega(\varphi_0, \psi_0)$  is a bounded weak solution of the stationary problem

$$\left\{ \begin{array}{ll} -\Delta u_i + f_i(x, u_1, u_2) = 0 & \text{in } \Omega \\ u_i = 0 & \text{on } \partial\Omega \end{array} \right.$$

**Proof :** From (3.3) we obtain  $\omega(\varphi_0, \psi_0) \neq \emptyset$ . Setting  $w_i = \lim_{n \rightarrow \infty} u_i(., t_n)$  and  $w = (w_1, w_2) \in \omega(\varphi_0, \psi_0)$ , we get that  $w = (w_1, w_2)$  is a solution of the Dirichlet problem for elliptic system. The proof is analogous to El Ouardi and de Th  lin [12] and is omitted here.

**Corollary 2.** Under the assumptions (H1) to (H10), we have

$$\mathcal{A} \subset (\mathcal{W}^{2,6}(\Omega))^2 \text{ if } N = 3$$

and

$$\mathcal{A} \subset (\mathcal{W}^{2,r}(\Omega))^2 \text{ for all } r < \infty \text{ if } N \leq 2.$$

Proof: Taking the inner product of (4.7) with  $u_i''$ , we get

$$\frac{d}{dt} \left( \sum_{i=1}^2 \|u_i'\|_{H_0^1(\Omega)}^2 \right) \leq c \left( \sum_{i=1}^2 |u_i'|_{H_0^1(\Omega)}^4 + \sum_{i=1}^2 |u_i'|_{L^2(\Omega)}^2 \right).$$

By uniform Gronwall's lemma, we get  $\sum_{i=1}^2 \|u_i'\|_{H_0^1(\Omega)}^2 \leq c, \forall t \geq T$ .

Then  $\sum_{i=1}^2 \|u_i'\|_{L^{\alpha_i}(\Omega)}^2 \leq c, \forall t \geq T$  for all  $t \geq \tau$  and  $\alpha_i = 6$  if  $N = 3$  or  $1 \leq \alpha_i < \infty$  if  $N \leq 2$ .

## 4. Dimension of the attractor $\mathcal{A}$

#### 4.1 Linearized problem

Let  $(\varphi_0, \psi_0) \in \mathcal{A}$ ; then by theorem3,  $u(t) = (u_1(t), u_2(t))$  belongs to a bounded subset of  $[H^2(\Omega)]^2$ . This fact allows us to linearize the system (S) along  $u(t)$ . Formally, the candidate for the linearized problem is

$$(S_L) \quad \begin{cases} U = (U_1, U_2) \in [L^2(0, T; H_0^1(\Omega))]^2 \\ \frac{\partial}{\partial t} (b'_i(u_i)U_i) - \Delta U_i + \sum_{j=1}^2 \frac{\partial f_i}{\partial u_j} U_j = 0 \\ U(0) = (U_1(0), U_2(0)) = U_0 \end{cases}$$

The existence and uniqueness of solution can be deduced from (4.0) below

$$U_i \in L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega)). \quad (4.0)$$

To deduce (4.0), Multiply the equation in  $(S_L)$  by  $b'_i(u_i)U_i$ , we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left( \sum_{i=1}^2 |b'_i(u_i)U_i|_{L^2(\Omega)}^2 \right) + \sum_{i=1}^2 (\nabla u_i, \nabla (b'_i(u_i)U_i))_{L^2(\Omega)} \\ &= \sum_{i=1}^2 \left( \sum_{j=1}^2 \frac{\partial f_i}{\partial u_j} U_j, b'_i(u_i)U_i \right)_{L^2(\Omega)} \end{aligned} \quad (4.1)$$

By the hypothesis on  $b_i$ , we have

$$\nabla (b'_i(u_i)U_i) = b'_i(u_i)\nabla U_i + b''_i(u_i)\nabla u_i \cdot U_i, \quad (4.2)$$

$$(\nabla U_i, b''_i(u_i)\nabla u_i \cdot U_i)_{L^2(\Omega)} \leq c |\nabla u_i|_{L^4(\Omega)} |U_i|_{L^4(\Omega)} |\nabla U_i|_{L^2(\Omega)}, \quad (4.3)$$

and

$$\sum_{i=1}^2 \left( \sum_{j=1}^2 \frac{\partial f_i}{\partial u_j} U_j, b'_i(u_i)U_i \right)_{L^2(\Omega)} \leq M \sum_{i=1}^2 \sum_{j=1}^2 |U_j U_i|_{L^1(\Omega)} \leq c \sum_{i=1}^2 |U_i|_{L^2(\Omega)}^2. \quad (4.4)$$

From (4.2) to (4.4), (4.1) becomes

$$\frac{1}{2} \frac{d}{dt} \left( \sum_{i=1}^2 |b'_i(u_i)U_i|_{L^2(\Omega)}^2 \right) + \gamma \sum_{i=1}^2 |U_i|_{H_0^1(\Omega)}^2 \leq c \sum_{i=1}^2 |U_i|_{L^2(\Omega)}^2 \leq c \sum_{i=1}^2 |b'_i(u_i)U_i|_{L^2(\Omega)}^2.$$

By standard application of Gronwall's inequality, we get (4.0).

#### 4.2 Differentiability of the Semigroup

We assume that  $f_i \in \mathcal{C}^2(R \times R)$  ( $\forall i = 1, 2$ ). Let  $u_0 = (\varphi_0, \psi_0)$ ,  $v_0 = (\overline{\varphi}_0, \overline{\psi}_0)$ ,  $S(t)$  be the solution of (S) and  $S'(t, u_0)$  the solution of  $(S_L)$ . The results of [6] imply the following proposition :

**Proposition 3.**

Assume (H1) to (H10), then for any  $(u_0, v_0) \in [L^\infty(\Omega) \times H^2(\Omega)]^2$ , we have  $|S(t)v_0 - S(t)u_0 - S'(t, u_0)(v_0 - u_0)|_{(L^2(\Omega))^2} \leq c(t)o(|v_0 - u_0|_{(L^2(\Omega))^2})$

We need the lemma 3 for the proof of proposition 3:

Let  $(u_1, u_2)$  and  $(v_1, v_2)$  be two solutions of (S) in  $\mathcal{A}$  with  $(u_1(0), u_2(0)) = (\varphi_0, \psi_0)$  and  $(v_1(0), v_2(0)) = (\varphi_1, \psi_1)$ . Setting  $w_1 = u_1 - v_1$  and  $w_2 = u_2 - v_2$ , we have

**Lemme 3** Assume (H1) to (H10). For all  $T > 0$ , there exists  $c(T) > 0$  such that for all  $t \in [0, T]$ ,

$$\sum_{i=1}^2 |w_i(t)|_{H_0^1(\Omega)}^2 \leq c(T) \sum_{i=1}^2 |w_i(0)|_{H_0^1(\Omega)}^2, \quad (4.7)$$

$$t \sum_{i=1}^2 |w_i(t)|_{H_0^1(\Omega)}^2 \leq c(T) \sum_{i=1}^2 |w_i(0)|_{L^2(\Omega)}^2, \quad (4.8)$$

and

$$\sum_{i=1}^2 |w_i'(t)|_{L^2(0,T;L^2(\Omega))}^2 \leq c(T) \sum_{i=1}^2 |w_i(0)|_{H_0^1(\Omega)}^2 \quad (4.9)$$

**Proof :** We have

$$\begin{cases} \frac{\partial b_i(u_i)}{\partial t} - \Delta u_i + f_i(x, u) = 0 \\ u(0) = (u_1(0), u_2(0)) = (\varphi_0, \psi_0) \end{cases} \quad \begin{cases} \frac{\partial b_i(v_i)}{\partial t} - \Delta v_i + f_i(x, v) = 0 \\ v(0) = (v_1(0), v_2(0)) = (\varphi_1, \psi_1) . \end{cases}$$

Thus, the difference  $w_i = u_i - v_i$  satisfies

$$b_i'(u_i)w_i' - \Delta w_i = [b_i'(v_i) - b_i'(u_i)]v_i' + f_i(v) - f_i(u). \quad (4.10)$$

Setting

$$F_{11} = \int_0^1 \frac{\partial f_1}{\partial u_1}(x, u_1 + \theta(u_2 - u_1), u_2) d\theta, F_{21} = \int_0^1 \frac{\partial f_1}{\partial u_2}(x, u_1, u_2 + \theta(u_2 - u_1)) d\theta, \\ F_{12} = \int_0^1 \frac{\partial f_2}{\partial u_1}(x, u_1 + \theta(u_2 - u_1), u_2) d\theta \text{ and } F_{22} = \int_0^1 \frac{\partial f_2}{\partial u_2}(x, u_1, u_2 + \theta(u_2 - u_1)) d\theta, \\ (4.10) \text{ becomes}$$

$$b_i'(u_i)w_i' - \Delta w_i = [b_i'(v_i) - b_i'(u_i)]v_i' + \sum_{j=1}^2 F_{ij}w_j. \quad (4.11)$$

We multiply (4.11) by  $\frac{w_i}{b'_i(u_i)}$

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \sum_{i=1}^2 |w_i|_{L^2(\Omega)}^2 + \sum_{i=1}^2 (\nabla w_i, \nabla (\frac{w_i}{b'_i(u_i)}))_{L^2(\Omega)} = \\ & \sum_{i=1}^2 (\frac{b'_i(v_i) - b'_i(u_i)}{b'_i(u_i)} v'_i, w_i)_{L^2(\Omega)} + \sum_{i=1}^2 \sum_{j=1}^2 (F_{ij} w_j, \frac{w_i}{b'_i(u_i)})_{L^2(\Omega)} \end{aligned} \quad (4.12)$$

And parallel to lemma 16 in [8], it is easy to see that (4.7), (4.8) and (4.9) hold.

**Proof of proposition 3 :** It is similar to the proof for the lemma 15 in [8, p.125] and is omitted.

### 4.3 Dimension Estimates

Consider the linearized problem

$$(S_L) \quad \begin{cases} U'_i = F'_i(u_i(\tau))U_i & \text{in } \Omega \times R^+ \\ U_i = 0 & \text{on } \partial\Omega \times R \\ U_i(0) = \xi_i \end{cases}$$

where  $F'_i(u_i(\tau))U_i = \frac{1}{b'_i(u_i(\tau))} \Delta U_i - \frac{b''_i(u_i(\tau))}{b'_i(u_i(\tau))} u'_i U_i - \frac{1}{b'_i(u_i(\tau))} \sum_{j=1}^2 \frac{\partial f_i}{\partial u_j} U_j$ .

This problem can be rewritten as

$$(L) \quad \begin{cases} U' = F'(u(\tau))U \\ U = 0 \\ U(0) = \xi \end{cases}$$

$$\text{where } U = \begin{pmatrix} U_1 \\ U_2 \end{pmatrix}, \quad F'(u(\tau)) = \begin{pmatrix} F'_1(u_1(\tau)) & 0 \\ 0 & F'_1(u_1(\tau)) \end{pmatrix}.$$

Let  $\bar{U}_1, \dots, \bar{U}_m$  be  $m$  solutions of (L) corresponding to the initial data  $\xi_1, \dots, \xi_m$  and  $\mathcal{Q}_m(\tau)$  be the orthogonal projector in  $H = L^2(\Omega) \times L^2(\Omega)$  such that  $\mathcal{Q}_m H \subset V = H_0^1(\Omega) \times H_0^1(\Omega)$ . If  $\{W^k = (w_1^k, w_2^k)\}_{k=1}^m$  is an orthonormal basis of  $\mathcal{Q}_m(\tau)H$ ; then

$$Tr(F'(u(\tau)) \circ \mathcal{Q}_m(\tau)) = \sum_{k=1}^m (F'(u(\tau))W^k, W^k)_H = \sum_{i=1}^2 \sum_{k=1}^m (F'_i(u_i(\tau))W_i^k, W_i^k)_{L^2(\Omega)}$$

and

$$\begin{aligned} & (F'_i(u_i(\tau))W_i^k, W_i^k)_{L^2(\Omega)} = (\Delta w_i^k, \frac{w_i^k}{b'_i(u_i(\tau))})_{L^2(\Omega)} - (\frac{b''_i(u_i(\tau))}{b'_i(u_i(\tau))} u'_i w_i^k, w_i^k)_{L^2(\Omega)} - \\ & (\frac{1}{b'_i(u_i(\tau))} \sum_{j=1}^2 \frac{\partial f_i}{\partial u_j} w_j^k, w_i^k)_{L^2(\Omega)}. \end{aligned}$$

Since  $u = (u_1, u_2) \in (L^\infty(0, +\infty, L^\infty(\Omega)))^2$ , we have  $b_i(u_i), b'_i(u_i)$ ,

$b_i''(u_i) \in (L^\infty(0, +\infty, L^\infty(\Omega)))^2$ , so

$$\sum_{i=1}^2 (F'_i(u_i(\tau))W_i^k, W_i^k)_{L^2(\Omega)} \leq \sum_{i=1}^2 (\Delta w_i^k, \frac{w_i^k}{b'_i(u_i(\tau))})_{L^2(\Omega)} + c \sum_{i=1}^2 \int_{\Omega} |u'_i| |w_i^k|^2 dx +$$

$$\int_{\Omega} \sum_{i=1}^2 \sum_{j=1}^2 \frac{\partial f_i(x, u)}{\partial u_j} w_j^k w_i^k dx. \quad (4.13)$$

Now, we have

$$\sum_{i=1}^2 (\Delta w_i^k, \frac{w_i^k}{b'_i(u_i(\tau))})_{L^2(\Omega)} \leq -c \sum_{i=1}^2 \|w_i^k\|_{H_0^1(\Omega)}^2 + cJ$$

where

$$\begin{aligned} J &= \sum_{i=1}^2 \int_{\Omega} |\nabla w_i^k| |w_i^k| |\nabla u_i| dx \leq c \left( \sum_{i=1}^2 \|w_i^k\|_{H_0^1(\Omega)}^2 \right)^{\frac{1}{2}} \left( \int_{\Omega} \sum_{i=1}^2 |w_i^k|^2 dx \right)^{\frac{1}{2}} \\ &\leq \frac{c}{2} \sum_{i=1}^2 \|w_i^k\|_{H_0^1(\Omega)}^2 + c \left( \int_{\Omega} \sum_{i=1}^2 |w_i^k|^2 dx \right) \end{aligned} \quad (4.14)$$

and

$$\int_{\Omega} \sum_{i=1}^2 \sum_{j=1}^2 \frac{\partial f_i(u)}{\partial u_j} w_j^k w_i^k dx \leq c \left( \int_{\Omega} \sum_{i=1}^2 |w_i^k|^2 dx \right). \quad (4.15)$$

According to (4.14) and (4.15), relation (4.13) becomes

$$\begin{aligned} \sum_{i=1}^2 (F'_i(u_i(\tau))W_i^k, W_i^k)_{L^2(\Omega)} &\leq -c \sum_{i=1}^2 \|w_i^k\|_{H_0^1(\Omega)}^2 + c \sum_{i=1}^2 \|w_i^k\|_{H_0^1(\Omega)}^2 + \\ &c \left( \int_{\Omega} \sum_{i=1}^2 |w_i^k|^2 dx \right) + c \sum_{i=1}^2 \int_{\Omega} |u'_i| |w_i^k|^2 dx \end{aligned} \quad (4.16)$$

Which leads to

$$\begin{aligned} Tr(F'(u(\tau)) \circ \mathcal{Q}_m(\tau)) &\leq -c_{105} \sum_{i=1}^2 \sum_{k=1}^m \|w_i^k\|_{H_0^1(\Omega)}^2 + c \sum_{i=1}^2 \sum_{k=1}^m \|w_i^k\|_{H_0^1(\Omega)}^2 + \\ &c \left( \int_{\Omega} \sum_{i=1}^2 \sum_{k=1}^m |w_i^k|^2 dx \right) + c \sum_{i=1}^2 \sum_{k=1}^m \int_{\Omega} |u'_i| |w_i^k|^2 dx. \end{aligned} \quad (4.17)$$

We set  $\rho(x) = \sum_{k=1}^m |w^k(x)|^2 = \sum_{i=1}^2 \sum_{k=1}^m |w_i^k(x)|^2$  and  $\gamma(t) = \max(|u'_1(t)|, |u'_2(t)|)$ ,  $\theta(t) = \int_{\Omega} \gamma(t)^{\frac{5}{2}} dx$ . So, we get

$$\sum_{i=1}^2 \sum_{k=1}^m \int_{\Omega} |u'_i| |w_i^k|^2 dx \leq \int_{\Omega} \gamma(t) \rho(x) dx \leq c \left[ \int_{\Omega} \gamma(t)^{\frac{5}{2}} dx \right]^{\frac{2}{5}} \left[ \int_{\Omega} \rho^{\frac{5}{3}} dx \right]^{\frac{3}{5}}. \quad (4.18)$$

Therefore by theorem 4.1 in [22], there exists  $c'_1 > 0$  such that :

$$\int_{\Omega} \rho^{\frac{5}{3}} dx \leq c'_1 \sum_{k=1}^m \|w^k(x)\|_{H_0^1(\Omega)}^2. \quad (4.19)$$

So, we get

$$\sum_{i=1}^2 \sum_{k=1}^m \int_{\Omega} |u'_i| |w_i^k|^2 dx \leq c \int_{\Omega} \gamma(t)^{\frac{5}{2}} dx + \frac{c}{5} \sum_{k=1}^m \|w^k(x)\|_{H_0^1(\Omega)}^2.$$

From (4.18) to (4.19), (4.16) becomes

$$Tr(F'(u(\tau)) \circ \mathcal{Q}_m(\tau) \leq -c \sum_{k=1}^m \|w^k\|_{H_0^1(\Omega)}^2 + c \int_{\Omega} \rho dx + c \int_{\Omega} \gamma(t)^{\frac{5}{2}} dx,$$

and as in [22], we obtain :

$$Tr(F'(u(\tau)) \circ \mathcal{Q}_m(\tau) \leq -cm^{1+\frac{2}{N}} + c'm + \theta(t). \quad (4.20)$$

$$\text{Setting } q_m(t) = \sup_{u_0 \in \mathcal{A}} \sup_{\substack{\xi_i \in H, \quad |\xi_i| \leq 1 \\ i=1, \dots, m}} \left\{ \frac{1}{t} \int_0^t Tr(F'(S(\tau)u_0)) \circ \mathcal{Q}_m(\tau) d\tau \right\}$$

and

$$q_m = \lim_{t \rightarrow +\infty} \sup q_m(t).$$

Then, by lemma 15 in [8, p.119], we have  $\int_0^\eta \theta d\tau \leq c(\eta)$  and  $u'_i \in L^\infty(\eta, +\infty, L^\infty(\Omega))$ .

Thus,  $q_m(t) \leq \frac{c(\eta)}{t} - cm^{1+\frac{2}{N}} + c'm + c'(\eta)$  and  $q_m \leq -cm^{1+\frac{2}{N}} + c'm + c'(\eta)$

and for all integers  $j > 0$ , we get  $\mu_1 + \mu_2 + \dots + \mu_j \leq q_j \leq -cj^{1+\frac{2}{N}} + c'j + c'(\eta)$ .

Hence

$$\mu_1 + \mu_2 + \dots + \mu_m < 0 \quad \text{for any } m < c''. \quad (4.21)$$

This shows that the fractal dimension of the attractor  $\mathcal{A}$  is finite and arguing as for theorem V3.3 in [22], we conclude to the following :

**Theorem 4.** Assume (H1) to (H10) and let  $m$  be an integer satisfying (4.21). Then

- ( i )  $\dim_{Fractale}(\mathcal{A}) \leq 2m$
- ( ii )  $\dim_H(\mathcal{A}) \leq m.$

**Acknowledgement.** The authors would like to thank the referee for his refereeing and helpful comments.

### References

- [1] ALT , H .W. and LUCKHAUSS, S . , Quasilinear elliptic and parabolic differential equations. Math.Z,183 (1983), pp311-341.
- [2] BAMBERGER , A. , Etude d'une équation doublement non linéaire, J.Func.Ana., 24 (77), pp148-155.
- [3] BLANCHARD , D. and FRANCFORT ,G., Study of doubly nonlinear heat equation with no growth assumptions on the parabolic term. Siam.J.Anal.Math, vol 9, n°5, sept. 1988.
- [4] DIAZ , J. and de THELIN , F. , On a nonlinear parabolic problem arising in some models related to turbulent flows, Siam.J.Anal.Math, vol25 , n°4, p1085-1111.
- [5] DIAZ , J. and J.F.Padial ,J. F. . Uniqueness and existence of solutions in the  $BV_t(Q)$  space to a doubly nonlinear parabolic problem.Publicacions Matemàtiques, vol40 (1996) , pp527-560.
- [6] DUNG.L. , Ultimately Uniform Boundedness of Solutions and Gradients for Degenerate Parabolic Systems. Nonlinear Analysis T.M.A.1998, in press.
- [7] DUNG.L. , Global attractors for a class of degenerate nonlinear parabolic systems, to appear in J.I.D.E.
- [8] EDEN , A. , MICHAUX , B. and RAKOTOSON , J.M.  
Doubly nonlinear parabolic type equations as dynamical systems, Journal of Dynamics and Differential Equations, vol3, n°1, 1991.
- [9] EDEN , A. , MICHAUX , B. and RAKOTOSON, J.M. ,  
Semi-discretized nonlinear evolution equations as discrete dynamical systems and error analysis, Indiana Uni.Math.Journal vol 39, N°3 (1990), p737-783.
- [10] EL HACHIMI , A. and de THELIN ,F. , Supersolutions and stabilisation of the solutions of the equation  $u_t - \Delta_p u = f(x, u)$ , *Part I*. Nonlinear Analysis T.M.A12, n°(88), p1385-1398 .
- [11] EL HACHIMI , A. and de THELIN ,F., Supersolutions and stabilisation of the solutions of the equation  $u_t - \Delta_p u = f(x, u)$ , *Part II*. Publicaciones Matemáticas, vol 35(1991) , pp347-362 .
- [12] EL OUARDI , H. and de THELIN, F., Supersolutions and Stabilization of the Solutions of a Nonlinear Parabolic System. Publicacions Matemàtiques, vol 33 (1989), 369-381.
- [13] FOIAS , C. , CONSTANTIN , P. and TEMAM , R. , Attractors representing turbulent flows. AMS Memoirs 53, No.314.
- [14] FOIAS , C. and TEMAM , R. , Structure of the set of stationary solutions of the Navier-Stokes equations. Comm. Pure Appl.Math.30.149-164.

- [15] LADYZHENSKAYA, O., and all , Linear and quasilinear equations of parabolic type, Trans.Amer. Math.Soc, providence, RI, 1968.
- [16] LANGLAIS, M. and PHILLIPS, D., Stabilization of solution of non-linear and degenerate evolution equations. Nonlinear Analysis,TMA, 9, (1985), 321-333.
- [17] LIONS , J. L. , Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod, Paris, 1969 .
- [18] Marion.M, Attractors for reaction-diffusion equation: existence and estimate of their dimension, Applicable Analysis (25) (1987), 101-147.
- [19] RAVIART , P.A. , Sur la résolution de certaines équations paraboliques non linéaires .Journ of Functional Analysis 5, 299-328 (1970).
- [20] SCHATZMAN , M. , Stationary solutions and asymptotic behavior of a quasilinear parabolic equation. Ind.Univ.J. 33 (1) ,1-29.
- [21] SIMON , J. , Régularité de la solution d'un problème aux limites non linéaire. Annales Fac. Sc. Toulouse3, série 5 (1981), p247-274.
- [22] TEMAM ,R. , Infinite dimensional dynamical systems in mechanics and physics.Applied Mathematical Sciences, n°68, springer-verlag (1988).
- [23] TSUTSUMI , M. , On solutions of some doubly nonlinear degenerate parabolic systems with absorption. .Journ.Math.Anal. and Applic.,132 (187-212) , 1988.

A.El Hachimi: UFR Mathématiques Appliquées et Industrielles  
 Faculté des Sciences, B.P 20, El Jadida - Maroc  
 e-mail adress: aelhachi@yahoo.fr  
 H. El Ouardi: UFR Mathématiques Appliquées et Industrielles  
 Faculté des Sciences, B.P 20, El Jadida - Maroc  
 and  
 Ecole Nationale Supérieure d'Electricité et de Mécanique  
 B.P 8118-Casablanca-Oasis - Maroc  
 e\_mail adress: h.elouardi@ensem-uh2c.ac.ma